NOW



# **PCT**

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

REC'D	07	MAR	2001

WIPO

PCT

## (PCT Article 36 and Rule 70)

Applicant's or agent's file reference	FOR FURTHER ACTIO		on of Transmittal of International xamination Report (Form PCT/IPEA/416)
342312003540			<u> </u>
International application No.	International filing date (day/	/month/year)	Priority date (day/month/year)
PCT/US99/29927	15 December 1999 (15.12.19		16 December 1998 (16.12.1998)
International Patent Classification (IPC)	or national classification and IF	PC	
IPC(7): C07K 7/56; A61K 38/12; A61P	31/10, 33/00 and US Cl.: 514/	/9; 530/317	
Applicant			
ELI LILLY AND COMPANY		<u> </u>	
Examining Authority and	is transmitted to the applicar	nt according to A	
2. This REPORT consists of	a total of $\frac{2}{3}$ sheets, include	ing this cover she	et.
which have been ame before this Authority	ended and are the basis for the (see Rule 70.16 and Section	his report and/or	description, claims and/or drawings sheets containing rectifications made nistrative Instructions under the PCT).
These annexes consist of a	total of <u>O</u> sheets.		
3. This report contains indica	ations relating to the following	ng items:	
I Basis of the rep	ort		
II Priority			
III Non-establishm	ent of report with regard to	novelty, inventive	e step and industrial applicability
IV Lack of unity o	f invention		
	nent under Article 35(2) wit tations and explanations sup		ty, inventive step or industrial ment
VI Certain docume	ents cited		
VII Certain defects	in the international applicati	ion	
VIII Certain observa	ations on the international ag	pplication	
Date of submission of the demand		Date of completion	of this report
14 July 2000 (14.07.2000)	2	9 January 2001 (29	.01.2001)
Name and mailing address of the IPEA/ Commissioner of Patents and Tradema Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	rks	Authorized officer Mukund Shah Telephone No. (703	Faurence for

Form PCT/IPEA/409 (cover sheet)(July 1998)

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

Internation application No.
PCT/US99/29927

I.	Basi	s of the report
1.	With	regard to the elements of the international application:*
	$\boxtimes$	the international application as originally filed.
	$\overline{\boxtimes}$	the description:
		pages 1-24 as originally filed
		pages NONE , filed with the demand
		pages NONE , filed with the letter of
	$\bowtie$	the claims:
		pages 25-36, as originally filed
		pages NONE , as amended (together with any statement) under Article 19
		pages NONE , filed with the demand pages NONE , filed with the letter of
	$\boxtimes$	the drawings.
		pages NONE , as originally filed
		pages NONE, filed with the demand  pages NONE, filed with the letter of
	$\triangle$	the sequence listing part of the description:
		pages NONE , as originally filed pages NONE , filed with the demand
		pages NONE , filed with the letter of
2	Wit	h regard to the language, all the elements marked above were available or furnished to this Authority in the
۷.	lang	uage in which the international application was filed, unless otherwise indicated under this item.
	The	se elements were available or furnished to this Authority in the following language which is:
		the language of a translation furnished for the purposes of international search (under Rule23.1(b)).
	$\Box$	the language of publication of the international application (under Rule 48.3(b)).
	H	the language of the translation furnished for the purposes of international preliminary examination (under Rules
	ш	55.2 and/or 55.3).
3.	Wit	h regard to any nucleotide and/or amino acid sequence disclosed in the international application, the
	inte	mational preliminary examination was carried out on the basis of the sequence listing:
		contained in the international application in printed form.
		filed together with the international application in computer readable form.
		furnished subsequently to this Authority in written form.
		furnished subsequently to this Authority in computer readable form.
		The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the
		international application as filed has been furnished.
		The statement that the information recorded in computer readable form is identical to the written sequence listing
		has been furnished.
4.	$\boxtimes$	The amendments have resulted in the cancellation of:
		the description, pages NONE
		the claims, Nos. NONE
		the drawings, sheets/fig NONE
5.	لبيا	This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**
th	is rep	ncement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in ort as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17). replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.
·	ınıy	reprocessing assess committing and assessment that are superior to make a superior and assessment to the superior and assess

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

Form PCT/IPEA/409 (Box V) (July 1998)

-		
Internati	application No.	
DCT/HS00	120027	

	PCT/US99/29927	
V. Reasoned statement under Article 35(2) citati ns and explanations supporting suc	with regard to novelty, inventive step or inc th statement	lustrial applicability;
1. STATEMENT		
Novelty (N)	Claims 1-23 Claims NONE	YES NO
Inventive Step (IS)	Claims 1-23 Claims NONE	YES NO
Industrial Applicability (IA)	Claims 1-23 Claims NONE	YES NO
2. CITATIONS AND EXPLANATIONS (Ru Claims 1-23 meet the criteria set out in PCT Article compositions or method of use of the compounds of	33(2)-(3), because the prior art does not teach or structural formula I.	airly suggest the compounds,
Claims 1-23 meet the criteria set out in PCT Article	33(4), for industrial applicability	
NONE		
	·	
÷		



# **PCT**

### INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 342312003540			of International Search Report s, where applicable, item 5 below.
International application No.	International filing date (day/mon	th/year) (Earliest) F	Priority Date (day/month/year)
PCT/US 99/29927	15/12/1999		16/12/1998
Applicant			
ELI LILLY AND COMPANY et	al	Annual Control	
This International Search Report has been according to Article 18. A copy is being to	n prepared by this International Seansmitted to the International Burea	arching Authority and is to au.	ransmitted to the applicant
This International Search Report consists  It is also accompanied by	of a total of sl a copy of each prior art document	neets. cited in this report.	
Basis of the report			
With regard to the language, the language in which it was filed, unit	international search was carried ou less otherwise indicated under this	it on the basis of the intellitem.	mational application in the
the international search w Authority (Rule 23.1(b)).	as carried out on the basis of a tra	nslation of the internation	al application fumished to this
b. With regard to any nucleotide an was carried out on the basis of th	e sequence listing :	sed in the international ap	pplication, the international search
I 😕	onal application in written form.	andable form	
	mational application in computer repetition in the state of the state	Badabie ioiiii.	
	this Authority in computer readble	form	
the statement that the sul	osequently furnished written sequents filed has been furnished.		eyond the disclosure in the
		dable form is identical to	the written sequence listing has been
2. X Certain claims were fou	nd unsearchable (See Box I).		
3. Unity of invention is lac	king (see Box II).		
4. With regard to the <b>title</b> ,			
the text is approved as su	ibmitted by the applicant.		
	shed by this Authority to read as fol		I MALT
CYCLIC PEPTIDE ANTIFU	NGAL AGENTS HAVING A	SUGAR SUBSTITU	ENT
5. With regard to the abstract,			
I ''	ibmitted by the applicant.		
the text has been establis within one month from th	shed, according to Rul 38.2(b), by date of mailing of this internations	this Authority as it appea al search report, submit c	ars in Box III. The applicant may, comments to this Authority.
6. The figure of the drawing to be pub	lished with the abstract is Figure N	o.	<del></del>
as suggested by th appl	icant.		Non of th figures.
because th applicant fai	led to suggest a figure.	·	
because this figure better	characteriz s th invention.		

### INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/29927

Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Inte	emational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claims 8-23 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
з. 🗌	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
This Inte	ernational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	The additional search fees w re accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

PCT/US 99/29927

A. CLASSI IPC 7	FICATION OF SUBJECT MATTER C07K7/56 A61K38/12 A61P31	1/10 A61P33/00	
	o International Patent Classification (IPC) or to both national class SEARCHED	sification and IPC	
Minimum do	cumentation searched (classification system followed by classifi	ication symbols)	
IPC 7			
Documental	tion searched other than minimum documentation to the extent the	at such documents are included in the fields se	earched
Electronic d	ata base consulted during the international search (name of data	a base and, where practical, search terms used	)
C DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the	a relevent pagagage	Relevant to claim No.
Calegory	Citation of document, with indication, where appropriate, or the	- Tolevall passages	neievair to dain 140.
Y	EP 0 757 058 A (LILLY CO ELI) 5 February 1997 (1997-02-05) the whole document		1–23
P,Y	WO 99 43337 A (LILLY CO ELI ;JA ANDREW (US); RODRIGUEZ MICHAEL 2 September 1999 (1999-09-02) the whole document		1–23
Α	WO 97 17365 A (MERCK & CO INC ; JAMES M (US); BOUFFARD FRANCES 15 May 1997 (1997-05-15) the whole document 		
Furth	ner documents are listed in the continuation of box C.	X Patent family members are listed	in annex.
° Special ca	tegories of cited documents :	"T" leter document nublished after the inte	metional filing data
consid "E" earlier o	ent defining the general state of the art which is not ered to be of particular relevance locument but published on or after the international	"T" later document published after the inte or priority date and not in conflict with cited to understand the principle or the invention "X" document of particular relevance; the c	the application but sory underlying the
which	ate nt which may throw doubts on priority claim(s) or is cited to establish the publication date of another n or other special reason (as specified)	cannot be considered novel or cannot involve an inventive step when the do "Y" document of particular relevance; the c	be considered to cument is taken alone laimed invention
"O" docume other r	ent referring to an oral disclosure, use, exhibition or	cannot be considered to involve an inv document is combined with one or mo ments, such combination being obviou in the art.	re other such docu-
later th	nan the priority date claimed	"&" document member of the same patent	
	ectual completion of the international search  May 2000	Date of mailing of the international sea 16/05/2000	rch report
<del></del>	nailing address of the ISA  — European-Patent Office, P.B. 5818 Patentlaan 2	Authorized officer	
	NL – 2280 HV Rijswijk Tel. (431–70) 340–2040, Tx. 31 651 epo nl, Fey: (431–70) 340–3016	Groenendijk, M	

## INT NATIONAL SEARCH REPORT

ormation on patent family members

rnational Application No PCT/US 99/29927

	atent document d in search repor	t	Publication date	1	Patent family member(s)	Publication date
EP	0757058	A	05-02-1997	US AU JP WO	5629289 A 6593896 A 11510165 T 9705163 A	26-02-1997 07-09-1999
WO	9943337	Α	02-09-1999	AU EP	2686099 A 0943623 A	
MO	9717365	Α	15-05-1997	AU AU EP HR JP	707664 B 7554096 A 0862579 A 960521 A 2000501076 T	29-05-1997 09-09-1998 30-04-1998

inte. onal Application No PCT/US 99/29927

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 CO7K7/56 A61k A61K38/12 A61P31/10 A61P33/00 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07K A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. Y EP 0 757 058 A (LILLY CO ELI) 1-23 5 February 1997 (1997-02-05) the whole document P,Y WO 99 43337 A (LILLY CO ELI ; JAMISON JAMES 1-23 ANDREW (US); RODRIGUEZ MICHAEL JOHN (U) 2 September 1999 (1999-09-02) the whole document WO 97 17365 A (MERCK & CO INC ; BALKOVEC JAMES M (US); BOUFFARD FRANCES A (US); DR) 15 May 1997 (1997-05-15) A the whole document Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents ; To later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "A" document defining the general state of the art which is not considered to be of particular relevance earlier document but published on or after the International \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled "O" document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed in the art. "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 9 May 2000 16/05/2000 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel: (+31-70) 340-2040; Tx: 31-651 epo nl; Groenendijk, M Fax: (+31-70) 340-3016

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/29927

Box I Observations whir certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 8-23 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
Claims Nos.:     because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
B x ii Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark in Protest  The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.



•

inte. onal Application No PCT/US 99/29927

	atent document I in search repor	t	Publication date		Patent family member(s)		Publication date
EP	0757058	A	05-02-1997	US AU JP WO		A A T A	13-05-1997 26-02-1997 07-09-1999 13-02-1997
WO	9943337	Α	02-09-1999	AU EP	2686099 0943623	A A	15-09-1999 22-09-1999
WO	9717365	A	15-05-1997	AU AU EP HR JP	0862579	B A A A T	15-07-1999 29-05-1997 09-09-1998 30-04-1998 02-02-2000

### **PCT**

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:

A1

(11) International Publication Number:

WO 00/35945

C07K 7/56, A61K 38/12, A61P 31/10,

33/00

\*\* | *|* 

(43) International Publication Date:

22 June 2000 (22.06.00)

(21) International Application Number:

PCT/US99/29927

(22) International Filing Date:

15 December 1999 (15.12.99)

(30) Priority Data:

60/112,433

16 December 1998 (16.12.98) US

(74) Agent: LEHNHARDT, Susan, K.; Morrison & Foerster LLP, 1290 Avenue of the Americas, New York, NY 10104-0012 (US).

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application

US

60/098,267 (CON)

Filed on

Not furnished

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RODRIGUEZ, John, Michael [US/US]; 1825 Sailing Court, Indianapolis, IN 46260 (US). NESLER, Michael, John [US/US]; 195 Lake Road, Bozrah, CT 06334 (US). ZWEIFEL, Mark, James [US/US]; 7519 Higdon Court, Indianapolis, IN 46214 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### **Published**

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: CYCLIC PEPTIDE ANTIFUNGAL AGENTS HAVING A SUGAR SUBSTITUENT

(57) Abstract

A compound represented by structure (I) is described where R<sup>8</sup> represents a sugar moiety. The compounds are shown to be useful as an antifungal and antiparasitic agent or as an intermediate to such an agent. Methods of treatment and pharmaceutical formulations containing compounds represented by structure (I) are also described.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		5256
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	_Liberia	SG	Singapore		
					<b>3.</b>		

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

5070

### CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



## 

(43) International Publication Date 22 June 2000 (22.06.2000)

**PCT** 

# (10) International Publication Number WO 00/35945 A1

- (51) International Patent Classification7: C07K 7/56, A61K 38/12, A61P 31/10, 33/00
- (21) International Application Number: PCT/US99/29927
- (22) International Filing Date:

15 December 1999 (15.12.1999)

(25) Filing Language:

**English** 

(26) Publication Language:

English

(30) Priority Data:

60/112,433 16 December 1998 (16.12.1998) U

- (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): RODRIGUEZ, John, Michael [US/US]; 1825 Sailing Court, Indianapolis, IN 46260 (US). NESLER, Michael, John [US/US]; 195 Lake Road, Bozrah, CT 06334 (US). ZWEIFEL, Mark, James [US/US]; 7519 Higdon Court, Indianapolis, IN 46214 (US).
- (74) Agent: JOHNSTON, Madeline, L.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).

- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:


- With international search report.
- (48) Date of publication of this corrected version:

5 July 2001

(15) Information about Correction: see PCT Gazette No. 27/2001 of 5 July 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CYCLIC PEPTIDE ANTIFUNGAL AGENTS HAVING A SUGAR SUBSTITUENT



(57) Abstract: A compound represented by structure (I) is described where R<sup>8</sup> represents a sugar moiety. The compounds are shown to be useful as an antifungal and antiparasitic agent or as an intermediate to such an agent. Methods of treatment and pharmaceutical formulations containing compounds represented by structure (I) are also described.



## **CORRECTED VERSION**

## (19) World Intellectual Property Organization International Bureau



## 

## (43) International Publication Date 22 June 2000 (22.06.2000)

### **PCT**

# (10) International Publication Number WO 00/35945 A1

- (51) International Patent Classification<sup>7</sup>:
  A61K 38/12, A61P 31/10, 33/00
- C07K 7/56,
- (21) International Application Number: PCT/US99/29927
- (22) International Filing Date:
  - 15 December 1999 (15.12.1999)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/112,433

16 December 1998 (16.12.1998) US

- (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): RODRIGUEZ, John, Michael [US/US]; 1825 Sailing Court, Indianapolis, IN 46260 (US). NESLER, Michael, John [US/US]; 195 Lake Road, Bozrah, CT 06334 (US). ZWEIFEL, Mark, James [US/US]; 7519 Higdon Court, Indianapolis, IN 46214 (US).
- (74) Agent: JOHNSTON, Madeline, I.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).

- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- With international search report.
- (48) Date of publication of this corrected version:


5 July 2001

(15) Information about Correction:

see PCT Gazette No. 27/2001 of 5 July 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CYCLIC PEPTIDE ANTIFUNGAL AGENTS HAVING A SUGAR SUBSTITUENT



- 0/35945 A
- (57) Abstract: A compound represented by structure (I) is described where R<sup>8</sup> represents a sugar moiety. The compounds are shown to be useful as an antifungal and antiparasitic agent or as an intermediate to such an agent. Methods of treatment and pharmaceutical formulations containing compounds represented by structure (I) are also described.



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7: C07K 7/56, A61K 38/12, A61P 31/10,

(11) International Publication Number:

WO 00/35945

33/00

(43) International Publication Date:

22 June 2000 (22.06.00)

(21) International Application Number:

PCT/US99/29927

A1

(22) International Filing Date:

15 December 1999 (15.12.99)

(30) Priority Data:

60/112,433

16 December 1998 (16.12.98)

US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application

US

60/098.267 (CON)

Filed on

Not furnished

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RODRIGUEZ, John, Michael [US/US]; 1825 Sailing Court, Indianapolis, IN 46260 (US). NESLER, Michael, John [US/US]; 195 Lake Road, Bozrah, CT 06334 (US). ZWEIFEL, Mark, James [US/US]: 7519 Higdon Court, Indianapolis, IN 46214 (US).

(74) Agent: LEHNHARDT, Susan, K.; Morrison & Foerster LLP, 1290 Avenue of the Americas, New York, NY 10104-0012 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, MIL, MR, NE, SN, TD, TG).

#### **Published**

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: CYCLIC PEPTIDE ANTIFUNGAL AGENTS HAVING A SUGAR SUBSTITUENT

### (57) Abstract

A compound represented by structure (I) is described where R8 represents a sugar moiety. The compounds are shown to be useful as an antifungal and antiparasitic agent or as an intermediate to such an agent. Methods of treatment and pharmaceutical formulations containing compounds represented by structure (I) are also described.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania .	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	Fi	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	T	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	ΥU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	u	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

### CYCLIC PEPTIDE ANTIFUNGAL AGENTS HAVING A SUGAR SUBSTITUENT

### TECHNICAL FIELD

The invention relates to anti-fungal/anti-parasitic agents. in particular, derivatives of Echinocandin compounds and their use in treatment of fungal and parasitic infections.

5

10

15

20

### BACKGROUND ART

A number of naturally occurring cyclic peptides are known in the art including Echinocandin B (A30912A), Aculeacin, Mulundocandin, Sporiofungin, L-671,329, and S31794/F1. In general, these cyclic peptides can be structurally characterized as a cyclic hexapeptide core (or nucleus) with an acylated amino group on one of the core amino acids. This acyl group is typically a fatty acid moiety forming a side chain off the nucleus. For example, Echinocandin B has a linolcoyl side chain while Aculeacin has a palmitoyl side chain.

These natural products have limited inherent antifungal and antiparasitic properties. The natural compounds can be structurally modified to enhance these properties or improve the compound's stability and/or water solubility. Turner et al., Cur. Pharm. Des. 2:209 (1996). For example, the fatty acid side chain can be removed from the cyclic peptide core to provide an amino nucleus which can then be re-acylated to provide semi-synthetic compounds.

### DISCLOSURE OF THE INVENTION

A compound represented by the following structure I is provided.

where R is an alkyl group, an alkenyl group, an alkynyl group. an aryl group, or heteroaryl group; R<sup>1</sup> is independently -H, -OH or -O-Pg; R<sup>2</sup> is -H, -CH<sub>3</sub>, -NH<sub>2</sub>, or -NH-Pg; R<sup>3</sup> is -H, -CH<sub>3</sub>, -CH<sub>2</sub>CONH<sub>2</sub>, -CH<sub>2</sub>CONH-Pg, -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, or -CH<sub>2</sub>CH<sub>2</sub>NH-Pg; R<sup>4</sup> is -H, -OH, or -O-Pg; R<sup>5</sup> is -OH, -OSO<sub>3</sub>H. or -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, *p*-halophenyl, *p*-halophenoxy, *p*-nitrophenyl, *p*-nitrophenoxy, benzyl, benzyloxy, *p*-halobenzyl, *p*-halobenzyl, *y*-halobenzyl, or *p*-nitrobenzyloxy; R<sup>6</sup> is -H, -OH, or -OSO<sub>3</sub>H; R<sup>7</sup> is -H or -CH<sub>3</sub>; t is an integer from 2-7; R<sup>8</sup> is a sugar moiety of the formula

where R<sup>9</sup> is independently -H, -OH, -N<sub>3</sub>, -O-Pg, -NH<sub>2</sub>, -NH-Pg, or a second sugar moiety comprising one to three sugar units selected from the group consisting of

5

15

$$R^{9b}$$
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 

$$R^{9b} \longrightarrow R^{9a} \longrightarrow R$$

' and mixtures

thereof, where R<sup>9a</sup> is -H, -OH, -N<sub>3</sub>, -NH<sub>2</sub>, -O-Pg, or -NH-Pg, R<sup>0b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or -NH-Pg, R<sup>9c</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>N<sub>3</sub>, -CH<sub>2</sub>OSO<sub>3</sub>H, -CH<sub>2</sub>NH-Pg, -CH<sub>2</sub>O-Pg, -CO<sub>2</sub>H, or -CO<sub>2</sub>-Pg, where R<sup>a</sup> is as defined above, and so long as no more than one R<sup>9</sup> is represented by said second sugar moiety; Pg is a protecting group (i.e., -O-Pg is a hydroxy protecting group, -NH-Pg is an amino protecting group, -CH<sub>2</sub>CONH-Pg is an amido protecting group and -CO<sub>2</sub>-Pg

is a carboxy protecting group); and pharmaceutically acceptable salts, esters, hydrates or solvates thereof.

The invention encompasses a pharmaceutical formulation of one or more pharmaceutical carriers, diluents or excipients and a compound represented by structure I described above.

5

10

15

20

25

30

The invention further encompasses a method of inhibiting fungal and parasitic activity by administering an effective amount of a compound represented by structure I to a recipient in need of thereof.

"Alkyl" refers to a hydrocarbon radical of the general formula  $C_nH_{2n+1}$  containing from 1 to 30 carbon atoms unless otherwise indicated. The alkane radical can be straight, branched, cyclic, or multi-cyclic. The alkane radical can be substituted or unsubstituted. Similarly, the alkyl portion of an alkoxy group, alkylthio group or alkanoate have the same definition as above.

"C1-C12 alkyl" refers to a straight or branched saturated alkyl chain having from one to twelve carbon atoms. C1-C12 alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, t-butyl, pentyl, 5-methylpentyl, hexyl, heptyl, 3,3-dimethylheptyl, octyl, 2-methyl-octyl, nonyl, decyl, undecyl and dodecyl. "C1-C12 alkyl" includes "C1-C6 alkyl", "C1-C4 alkyl", and "C3-C12 cycloalkyl."

"C<sub>3</sub>-C<sub>12</sub> cycloalkyl" refers to a cyclic saturated alkyl chain having from 3 to 12 carbon atoms. Moreover, "C<sub>3</sub>-C<sub>12</sub> cycloalkyl" includes "C<sub>3</sub>-C<sub>7</sub> cycloalkyl", *i.e.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

"C1-C12 alkoxy" refers to a C1-C12 alkyl group attached through an oxygen atom. C1-C12 alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, sec-butoxy, n-pentoxy, 5-methyl-hexoxy, heptoxy, octyloxy, decyloxy and dodecyloxy. "C1-C12 alkoxy" includes "C1-C6 alkoxy", "C3-C7 alkoxy", and "C1-C4 alkoxy".

"C1-C12 alkylthio" refers to a C1-C12 alkyl group attached through a sulfur atom. C1-C12 alkylthio groups include, but are not limited to, methylthio, ethylthio, propylthio, isopropylthio, butylthio, 3-methyl-heptylthio, octylthio, and 5,5-dimethyl-hexylthio. "C1-C12 alkylthio" includes "C1-C6 alkylthio" and "C1-C4 alkylthio."

"Alkenyl" refers to an acyclic hydrocarbon containing at least one carboncarbon double bond. The alkene radical can be straight, branched, cyclic, or multicyclic. The alkene radical can be substituted or unsubstituted.

"Alkynyl" refers to an acyclic hydrocarbon containing at least one carboncarbon triple bond. The alkyne radical can be straight, or branched. The alkyne radical can be substituted or unsubstituted.

5

10

15

20

25

30

"C2-C12 alkynyl" refers to a straight or branched mono-alkynyl chain having from two to twelve carbon atoms. C2-C12 alkynyl groups include, but are not limited to, ethynyl, 1-propyn-1-yl, 1-propyn-2-yl, 1-butyn-1-yl, 1-butyn-3-yl, 1-pentyn-3-yl, 4-pentyn-2-yl, 1-hexyn-3-yl, 3-hexyn-1-yl, 5-methyl-3-hexyn-1-yl, 5-octyn-1-yl, 7-octyn-1-yl, 4-decyn-1-yl and 6-decyn-1-yl.

"Aryl" refers to aromatic moieties having single (e.g., phenyl) or fused ring systems (e.g., naphthalene, anthracene, phenanthrene, etc.). The aryl groups can be substituted or unsubstituted. Substituted aryl groups include a chain of aromatic moieties (e.g., biphenyl, terphenyl, phenylnaphthalyl, etc.).

"Heteroaryl" refers to aromatic moieties containing at least one heteratom within the aromatic ring system (e.g., pyrrole, pyridine, indole, thiophene, furan, benzofuran, imidazole, pyrimidine, purine, benzimidazole, quinoline, etc.). The aromatic moiety can be a single or fused ring system. The heteroaryl groups can be substituted or unsubstituted.

Within the field of organic chemistry and particularly within the field of organic biochemistry, it is widely understood that significant substitution of compounds is tolerated or even useful. In the present invention, for example, the term alkyl group allows for substituents which are a classic alkyl, such as methyl, ethyl, propyl, n-butyl, i-butyl, hexyl, isooctyl, dodecyl, stearyl, etc. The term group specifically envisions and allows for substitutions on alkyls which are common in the art, such as hydroxy, halogen, alkoxy, carbonyl, keto, ester, carbamato, etc., as well as including the unsubstituted alkyl moiety. However, it is generally understood by those skilled in the art that the substituents should be selected so as to not adversely affect the pharmacological characteristics of the compound or adversely interfere with the use of the medicament. The same is true for each of the other groups (i.e., aryl, alkynyl, alkenyl, heteroaryl). Suitable substituents for any of the groups defined above include alkyl, alkenyl, alkynyl, aryl, halo, hydroxy, alkoxy, aryloxy, mercapto,

alkylthio, arylthio, mono- and di-alkyl amino, quaternary ammonium salts, aminoalkoxy, hydroxyalkylamino, aminoalkylthio, carbamyl, carbonyl, carboxy, glycolyl, glycyl, hydrazino, guanyl, and combinations thereof.

"Halo" refers to chloro, fluoro, bromo and iodo.

5

10

"O-Pg" and "hydroxy protecting group" refer to a substituent of a hydroxy group that is commonly employed to block or protect the hydroxy functionality while reactions are carried out on other functional groups on the compound. This substituent, when taken with the oxygen to which it is attached, can form an ether, e.g., methyl, methoxymethyl, and benzyloxymethyl ether, a silyl ether, an ester, e.g. acetoxy, or a sulfonate moiety, e.g. methane and p-toluenesulfonate. The exact genus and species of hydroxy protecting group is not critical so long as the derivatized hydroxy group is stable to the conditions of subsequent reaction(s) and the protecting group can be removed at the appropriate point without disrupting the remainder of the molecule. A preferred hydroxy protecting group is acetyl. Specific examples of hydroxy protecting groups are described in Greene, "Protective Groups in Organic Synthesis," John Wiley and Sons, New York, N.Y., (2nd ed., 1991), ("Greene") chapters 2 and 3 and in the Preparations and Examples sections which follow.

15

group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. When p is 0, the amino protecting group, when taken with the nitrogen to which it is attached, forms a cyclic imide, e.g., phthalimido and tetrachlorophthalimido. When p is 1, the protecting group, when taken with the nitrogen to which it is attached, can form a carbamate, e.g., methyl, ethyl, and 9-fluorenylmethylcarbamate; or an amide, e.g., N-formyl and N-

"NHp-Pg" and "amino protecting group" refer to a substituent of the amino

25

20

acetylamide. The exact genus and species of amino protecting group employed is not critical so long as the derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the intermediate molecule and the protecting group can be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other amino protecting group(s). Preferred amino protecting groups are t-butoxycarbonyl (t-Boc), allyloxycarbonyl, phthalimido, and

benzyloxycarbonyl (CbZ). Further examples of groups referred to by the above terms

30

are described in Greene at chapter 7.

5

10

15

20

25

30

"-CO<sub>2</sub>-Pg" and "carboxy protecting group" refer to a substituent of a carbonyl that is commonly employed to block or protect the carboxy functionality while reactions are carried out on other functional groups on the compound. This substituent, when taken with the carbonyl to which it is attached. can form an ester, e.g., C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, benzyl, substituted benzyl, benzhydryl, substituted benzhydryl, trityl, substituted trityl, and trialkylsilyl ester. The exact species of carboxy protecting group is not critical so long as the derivatized carboxy group is stable to the conditions of subsequent reaction(s) and the protecting group can be removed at the appropriate point without disrupting the remainder of the molecule. Other examples of groups referred to by the above terms are described in *Greene*, at chapter 5.

"C(O)NH-Pg" and "amido protecting group" refer to a substituent of an amide commonly employed to block or protect the amino portion while reacting other functional groups on the compound. This protecting group, when taken with the nitrogen to which it is attached, can form an amide, e.g. N-allyl, N-methoxymethyl, and N-benzyloxymethyl amide. The exact species of amido protecting group employed is not critical so long as the derivatized amido group is stable to the condition of subsequent reaction(s) on other positions of the intermediate molecule and the protecting group can be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other amido protecting group(s). Other examples of groups referred to by the above terms are described in *Greene*, at chapter 7, pg. 397.

"Carbonyl activating group" refers to a substituent of a carbonyl that promotes nucleophilic addition reactions at that carbonyl. Suitable activating substituents are those which have a net electron withdrawing effect on the carbonyl. Such groups include, but are not limited to, alkoxy, aryloxy, nitrogen containing aromatic heterocycles, or amino groups such as oxybenzotriazole, imidazolyl, nitrophenoxy, pentachlorophenoxy, N-oxysuccinimide, N,N'-dicyclohexylisoure-O-yl, N-hydroxy-N-methoxyamino; acetates, formates, sulfonates such as methanesulfonate, ethanesulfonate, benzenesulfonate, or p-tolylsulfonate; and halides such as chloride, bromide, or iodide.

"Pharmaceutical" or "pharmaceutically acceptable" means substantially non-toxic and substantially non-deleterious to the recipient. "Pharmaceutical formulation"

means that the carrier, solvent, excipients and salt are compatible with the active ingredient of the formulation (i.e., Compound I).

"Pharmaceutical salt" or "pharmaceutically acceptable salt" refers to salts of the compounds represented by structure I that are substantially non-toxic to the recipient at the doses administered. Typical pharmaceutical salts include those prepared by reaction of the compounds of the invention with a mineral or organic acid or inorganic base. Such salts are known as acid addition and base addition salts. For further exemplification, see *e.g.* Berge et al., *J. Pharm. Sci.*, 66:1 (1977).

5

10

15

20

25

30

"Solvate" is an aggregate that comprises one or more molecules of the solute, such as a formula I compound, with one or more molecules of a pharmaceutical solvent, including, but not limited to, water and ethanol. "Suitable solvent" is any solvent, or mixture thereof, inert to the ongoing reaction that sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction.

"Thermodynamic base" is a base that provides a reversible deprotonation of an acidic substrate or is a proton trap for protons produced as reaction byproducts, and is reactive enough to effect the desired reaction without significantly effecting any undesired reactions. Examples of thermodynamic bases include, but are not limited to, acetates, acetate dihydrates, carbonates, bicarbonates, C<sub>1</sub>-C<sub>4</sub> alkoxides, and hydroxides (e.g. silver, lithium, sodium, or potassium acetate, acetate dihydrate, carbonate, bicarbonate, methoxide, or hydroxide), tri-(C<sub>1</sub>-C<sub>4</sub> alkyl)amines, or aromatic nitrogen containing heterocycles (e.g. imidazole and pyridine).

"Inhibiting" includes prohibiting, stopping, retarding, alleviating, ameliorating, halting, restraining, slowing or reversing the progression, or reducing the severity of the growth or any attending characteristics, symptoms, and results from the existence of a parasite or fungus. These methods include both medical therapeutic (acute) and/or prophylactic (prevention) administration as appropriate.

"Effective amount" refers to an amount of a compound of formula I which is capable of inhibiting fungal and/or parasitic activity.

"Recipient" includes mammals, preferably, humans.

### BEST MODE FOR CARRYING OUT THE INVENTION

Compounds represented by structure I have now been found to be useful as antifungal and antiparasitic agents or as an intermediate thereof. The most convenient

5

10

15

20

means of producing compounds represented by structure I is by modifying naturally occurring compounds.

For illustrative purposes, Scheme I (below) starts with a specific Echinocandin derivative. However, one could begin with any natural product, semi-synthetic or synthetic Echinocandin-type compound containing a hemiaminal group.

The term "Echinocandin-type compounds" refers to compounds having the following general structure including any simple derivatives thereof:

wherein R is an alkyl group, an alkenyl group, an alkynyl group, an aryl group, or heteroaryl group; R<sup>1</sup> is -H or -OH; R<sup>2</sup> is -H, -NH<sub>2</sub> or -CH<sub>3</sub>; R<sup>3</sup> is -H, -CH<sub>3</sub>, -CH<sub>2</sub>CONH<sub>2</sub> or -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>; R<sup>4</sup> is -H or -OH; R<sup>5</sup> is -OH, -OSO<sub>3</sub>H, or -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, *p*-halophenyl, *p*-halophenoxy, *p*-nitrophenyl, *p*-nitrophenoxy, benzyl, benzyloxy, *p*-halobenzyl, *p*-halobenzyl, *p*-halobenzyloxy, *p*-nitrobenzyl, or *p*-nitrobenzyloxy; R<sup>6</sup> is -H, -OH, or -OSO<sub>3</sub>H; R<sup>7</sup> is -H or -CH<sub>3</sub>; and pharmaceutically acceptable salts, esters, hydrates or solvates thereof.

"Natural product" is those secondary metabolites, usually of relatively complex structure, which are of more restricted distribution and more characteristic of a specific source in nature. Suitable natural product starting materials of the Echinocandin cyclopeptide family include Echinocandin B, Echinocandin C, Aculeacin Ay, Mulundocandin, Sporiofungin A, Pneumocandin A<sub>0</sub>, WF11899A, and Pneumocandin B<sub>0</sub>.

The cyclic peptides used in the invention can be produced by culturing various microorganisms. In general, the cyclic peptides can be characterized as a cyclic hexapeptide nucleus with an acylated amino group on one of the amino acids. The

5

10

15

20

25

amino group on the naturally-occurring cyclic peptide is typically acylated with a fatty acid group forming a side chain off the nucleus. Examples of naturally-occurring acyl groups include, but are not limited to, linoleoyl (Echinocandin B, C and D), palmitoyl (Aculeacin Aγ and WF11899A), stearoyl, 12-methylmyristoyl (Mulundocandin), 10,12-dimethylmyristoyl (Sporiofungin A and Pneumocandin A<sub>0</sub>).

Semi-synthetic derivatives can be generally prepared by removing the fatty acid side chain from the cyclic peptide nucleus to produce a free amino group (i.e., no pendant acyl group -C(O)R). The free amine is then re-acylated with a suitable acyl group. For example, the echinocandin B nucleus has been re-acylated with certain nonnaturally occurring side chain moieties to provide a number of antifungal agents.

U.S. Patent No. 4,293,489. The N-acyl side chain encompasses a variety of side chain moieties known in the art. Suitable side chain moieties include substituted and unsubstituted alkyl groups, alkenyl groups, alkynyl groups, aryl groups, heteroaryl groups and combinations thereof. Preferably, the side chain contains both a linearly rigid section and a flexible alkyl section to maximize antifungal potency.

Representative examples of preferred acyl side chains include R groups having the following structures:

where A, B, C and D are independently hydrogen,  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkynyl,  $C_1$ - $C_{12}$  alkoxy,  $C_1$ - $C_{12}$  alkylthio, halo, or -O- $(CH_2)_m$ -[O- $(CH_2)_n$ ]<sub>p</sub>-O- $(C_1$ - $C_{12}$  alkyl) or -O- $(CH_2)_q$ -X-E; m is 2, 3 or 4; n is 2, 3 or 4; p is 0 or 1; q is 2, 3 or 4; X is pyrrolidino, piperidino or piperazino; and E is hydrogen,  $C_1$ - $C_{12}$  alkyl,  $C_3$ - $C_{12}$  cycloalkyl, benzyl or  $C_3$ - $C_{12}$  cycloalkylmethyl.

Scheme I illustrates the general semi-synthetic route described above where a natural product (Compound II(a)) is modified to provide an intermediate (Compound

II(d)) which is then further modified to provide a representative example of Compound I as illustrated in Scheme II.

Scheme I

5

Cyclic peptides represented by structure II(a) can be prepared by fermentation of known microorganisms. For example, the cyclic peptide II(a) where R<sup>1</sup> and R<sup>4</sup> are each hydroxy, R<sup>2</sup>, R<sup>3</sup> and R<sup>7</sup> are each methyl (cyclic nucleus corresponding to A-

II(d)

30912A) can be prepared using the procedure detailed in U.S. Patent No. 4,293,482. Cyclic peptide II(a) where R<sup>1</sup> is hydroxy, R<sup>2</sup>, R<sup>3</sup> and R<sup>7</sup> are each methyl, and R<sup>4</sup> is hydrogen (cyclic nucleus corresponding to A-30912B) can be prepared using the procedure detailed in U.S. Patent No. 4,299,763. Aculeacin can be prepared using the procedure detailed in U.S. Patent No. 3,978,210. Cyclic peptide II(a) where R<sup>3</sup> is CH<sub>2</sub>C(O)NH<sub>2</sub>, R<sup>7</sup> is methyl, R<sup>2</sup> is hydrogen, and R<sup>1</sup> and R<sup>4</sup> are hydroxy can be prepared using the procedure detailed in U.S. Patent No. 5,198,421.

5

10

15

20

25

30

Cyclic peptide II(a) can be deacylated using procedures known in the art to provide an amino nucleus represented by structure II(b). This reaction is typically carried out enzymatically by exposing the naturally occurring cyclic peptide to a deacylase enzyme. The deacylase enzyme can be obtained from the microorganism Actinoplanes utahensis and used substantially as described in U.S. Patent Nos. 4,293,482 and 4,304,716. The deacylase enzyme can also be obtained from the Pseudomonas species. Deacylation can be accomplished using whole cells of A. utahensis or Pseudomonas or the crude or purified enzyme thereof or using an immobilized form of the enzyme. See European Patent Application No. 0 460 882. Examples of naturally occurring cyclic peptides that can be used as starting materials include, but are not limited to, aculeacin (palmitoyl side chain), tetrahydroechinocandin B (stearoyl side chain), mulundocandin (branched C15 side chain), L-671,329 (C16 branched side chain), S 31794/F1 (tetradecanoyl side chain), sporiofungin (C<sub>15</sub> branched side chain), FR901379 (palmitoyl side chain). A preferred cyclic peptide is echinocandin B (Compound II(a) where R<sup>1</sup> and R<sup>4</sup> are each hydroxy, R<sup>2</sup>, R<sup>3</sup> and R<sup>7</sup> are each methyl, and R<sup>nat</sup> is linolecyl).

The amino nucleus II(b) can be re-acylated, by procedures taught in U.S. Patent Nos. 5,646,111, and 5,693,611 to provide compounds represented by structure II(c). See Preparation 12 below for an example of this transformation. Also see, U.S. Patent Nos. 5,646,111 and 5,693,611 for preparation of the acyl groups at R. Cyclic peptides II(c) where R contains 1 or more heterocyclic rings can be prepared as taught in U.S. Patent No. 5,693,611.

Compound II(d) can then be prepared by selective etherification of Compound II(c) as taught in *J. Antibiotics*, 51:239-242 (1998) and U.S. Patent No. 5,652,213.

The compounds represented by structure I can be prepared from compounds of structure II(d) as illustrated in Scheme II below where R<sup>10</sup> is a carbonyl activating group and t, R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>7</sup> and R<sup>9</sup> are as defined above.

Scheme II

5

10

15

Compound I can be prepared by adding either Compound III or IV to Compound II(a) dissolved or suspended in a suitable solvent in the presence of a suitable thermodynamic base. A convenient and preferred solvent for the reaction is dimethylformamide while a convenient and preferred base is triethylamine. The reaction can be performed at from 0°C to the reflux temperature of the mixture but is typically performed at ambient temperatures for about 24 hours. See Example 1 below for an example of specific reaction conditions.

Compounds represented by structures III and IV are known in the art and if not commercially available can be synthesized by techniques well known in synthetic chemical arts. Collins et al., "Monosaccharides: Their Chemistry and Their Roles in Natural Products," John Wiley and Sons, New York, NY, 1995; and "Methods in Carbohydrate Chemistry", Vol VI, Academic Press, New York, N.Y., 1980.

Compounds III and IV where R<sup>9</sup> is hydroxy are known as carbohydrates or monosaccharides (sugars). These sugars can be modified by replacing one or more hydroxy groups with hydrogen, azide, or amino to provide the other derivatives of Compounds III and IV including disaccharides and polysaccharides where R<sup>9</sup> is a second sugar moiety. Such compounds can be prepared as illustrated in Scheme 3 below where Lg is an activated hydroxy leaving group.

5

10

15

20

A commercially available Compound VI can have its hydroxy group(s) activated for nucleophilic displacement by standard techniques known in the art. For example, the hydroxy group can be sulfonylated with methane-, benzene-, or ptoluene-sulfonyl chloride (or bromide) to provide a Compound VII where Lg is OSO<sub>2</sub>Me, OSO<sub>2</sub>-phenyl, or OSO<sub>2</sub>-p-toluenyl. At this point, the leaving group can be displaced by azide ion, e.g., from sodium or potassium azide. Alternatively, the leaving group can be displaced by iodide ion from, e.g., sodium or potassium iodide. The resulting Compound VIII can be reduced to form a Compound IX where one or more of R<sup>9a</sup> or R<sup>9b</sup> is amino or hydrogen by catalytic hydrogenation or with a reducing agent such as nickel chloride hexahydrate. It is preferred that when an amino group is desired in the final product Compound I, that any azido groups are converted to amino groups after coupling to Compound II(d).

Compound I, where any of R<sup>9</sup> is amino can be formed from a Compound I where R<sup>9</sup> is azido as described by analogous procedures well known in the art. See, e.g., Larock, "Comprehensive Organic Transformations," pg. 409, VCH Publishers, New York, N.Y., 1989.

Compound I where R<sup>5</sup>, R<sup>9</sup>, R<sup>9a</sup>, R<sup>9b</sup>, and/or R<sup>9c</sup> is a hydroxy group, can be phosphorylated or phosphonylated by reaction with an appropriately substituted dichloro- phosphate or phosphonic acid of formula V

5

in the presence of a suitable base to provide, following an aqueous work-up, to produce Compound I where R<sup>5</sup>, R<sup>9</sup>, R<sup>9a</sup>, R<sup>9b</sup>, and/or R<sup>9c</sup> are moieties of the formula

10

Suitable bases include lithium trimethylsilanolate (LiOTMS), and lithium bis(trimethylsilyl)amide (LHMDS). A preferred solvent is an aprotic solvent such as tetrahydrofuran and/or dimethylformamide. U.S. Patent No. 5,693,611.

Alternatively, the compounds represented by structure I where R<sup>9b</sup> is hydroxy and/or R<sup>9c</sup> is hydroxymethyl can be sulfated by reaction with a suitable sulfation reagent. Guiseley et al., *J. Org. Chem.*, 26:1248 (1961). The protected compound of structure I can have its protecting group(s) removed to form a deprotected Compound I. Initial choices of protecting groups, and methods for their removal, are well known in the art. See, *e.g.*, *Greene*.

20

15

Pharmaceutical salts are typically formed by reacting Compound I with an equimolar or excess amount of acid or base. The reactants are generally combined in a mutual solvent such as diethylether, tetrahydrofuran, methanol, ethanol, isopropanol, benzene, and the like for acid addition salts, or water, an alcohol or a chlorinated solvent such as methylene chloride for base addition salts. The salts normally precipitate out of solution within about one hour to about ten days and can be isolated by filtration or other conventional methods.

25

Acids commonly employed to form acid addition salts are inorganic acids including, but not limited to, hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and organic acids such as p-toluenesulfonic, methanesulfonic acid, ethanesulfonic acid, oxalic acid, p-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, tartaric acid, benzoic acid, acetic acid.

Base addition salts include those derived from inorganic bases, including, but not limited to, ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates. Bases useful in preparing salts of this invention include, but are not limited to, sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide and calcium carbonate.

The particular counterion forming a part of any salt of this invention is not of a critical nature, provided the salt as a whole is pharmacologically acceptable and the counterion does not contribute undesired qualities to the salt as a whole. Preferred pharmaceutical acid addition salts are those formed with mineral acids such as hydrochloric acid and sulfuric acid, and those formed with organic acids such as maleic acid, tartaric acid, and methanesulfonic acid. Preferred pharmaceutical base addition salts are the potassium and sodium salt forms.

Preferred compounds of the present invention are those compounds represented by structure I where  $R^1$  is hydroxy at each occurrence;  $R^4$  is hydroxy;  $R^2$ ,  $R^3$ , and  $R^7$  are each methyl; R is a moiety of the formula

R<sup>a</sup> is methyl or methoxy; or a pharmaceutically acceptable salt or solvate thereof. More preferable are those compounds wherein R<sup>5</sup> is hydroxy; R is a moiety of the formula

R<sup>8</sup> is a moiety of the formula:

5

10

15

20

D is hydrogen or C<sub>3</sub>-C<sub>7</sub> alkoxy; R<sup>9</sup> is independently hydrogen, hydroxy, amino, or a moiety of the formula:

where R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or -NH-Pg and n is 1, 2, or 3; or a pharmaceutically acceptable salt or solvate thereof. Even more preferable are those compounds wherein D is n-pentoxy; R<sup>9</sup> is independently hydroxy or amino; or a pharmaceutically acceptable salt or solvate thereof. Most preferred are those compounds wherein R<sup>9</sup> is hydroxy at each occurrence; and t is 2; or a pharmaceutically acceptable salt or solvate thereof.

5

10

15

20

25

The optimal time for performing the reactions of Schemes 1 - 3 can be determined by monitoring the progress of the reaction by conventional chromatographic techniques. Choice of reaction solvent is generally not critical so long as the solvent employed is inert to the ongoing reaction and sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction. Unless otherwise indicated, all of the reactions described herein are preferably conducted under an inert atmosphere. A preferred inert atmosphere is nitrogen. Once a reaction is complete, the intermediate compound can be isolated by procedures well-known in the art, for example, the compound can be crystallized or precipitated and then collected by filtration, or the reaction solvent can be removed by extraction, evaporation or decantation. The intermediate compound can be further purified, if desired, by common techniques such as crystallization, precipitation, or chromatography over solid supports such as silica gel, alumina and the like, before carrying out the next step of the reaction scheme.

The following examples are meant to illustrate but not limit the invention. All references cited herein are hereby incorporated herein by reference.

The following Preparations and Examples further describe synthesis of the compounds. "Fast atom bombardment mass spectroscopy" and "high performance liquid chromatography" are abbreviated "MS(FAB)" and "HPLC", respectively.

### Preparation 1

The A-30912A nucleus (60.2 mmol) and the 2,4,5-trichlorophenol ester of [(4"-pentyloxy)-1,1':4',1"-terphenyl]-4-carboxylic acid (26.0 g, 48.2 mmol) were combined in 8.5 L of dimethylformamide. The resultant reaction mixture was stirred for approximately 48 hours at room temperature (RT) and the solvent was removed *in vacuo* to provide a residue. This residue was slurried in ether, collected by filtration, washed with methylene chloride and dissolved in methanol or a 1:1 (v/v) acetonitrile/water mixture. The resultant solution is subjected to reverse phase HPLC (C18; eluent of 20-40% aqueous acetonitrile containing 0.5% monobasic ammonium phosphate (w/v); 20 mL/min.; 230 nm). After removing the unreacted A30912A nucleus, the desired product is eluted from the column using an eluent of aqueous acetonitrile. The fractions containing the desired product are combined and then concentrated *in vacuo* or lyophilized to provide 18 g of the title compound.

MS(FAB): 1140.5103 (M<sup>+1</sup>).

5

10

15

### Preparation 2

The compound of Preparation 1 (1.08 g, 0.9 mmol) and ethanolamine hydrochloride (185 mg, 1.9 mmol) were dissolved in about 20 mL of dimethylsulfoxide. Hydrogen chloride was blown over the solution for about 5 seconds. The resulting mixture was stirred at room temperature under nitrogen for 28 hours. The reaction solution was subjected to reverse-phase HPLC (C18; eluent of 60% acetonitrile in water containing 0.1% trifluoroacetic acid (v/v); 40 mL/min.; 280 nm). The fractions containing the desired product were combined and concentrated in vacuo or lyophilized to provide the title compound. MS(FAB):1183.5577 (M+H).

### Examples 1 and 2

Examples 1 and 2 have the following base structure:

5

10

α-D-Galacturonic acid (388 mg, 2.0 mmol) and N-hydroxysuccinimide (NHS) (230 mg, 2.0 mmol) were dissolved in anhydrous dimethylformamide. The mixture was cooled in an ice bath under nitrogen and dicyclohexylcarbodiimide (DCC) (412 mg, 2.0 mmol) was added. The ice bath was removed and the resulting mixture was stirred overnight and then the reaction solution was filtered directly into a flask containing the compound of Preparation 2 (240 mg, 0.2 mmol). Triethylamine (0.030 mL, 0.2 mmol) was then added and the resulting mixture was stirred at room temperature under nitrogen for 48 hours. The resultant solution was subjected to reverse-phase HPLC (C18; eluent of 60% acetonitrile in water; 280 nm). The fractions containing the desired product were combined and then concentrated *in* vacuo or lyophilized to provide the title compound. MS(FAB): 1381 (M<sup>-</sup>+Na).

5

10

15

20

25

Example 2

OH
HO
HO
HO

D-Glucuronic acid (194 mg, 1.0 mmol) and the compound of Preparation 2 were converted to the title compound by the procedure of Example 1

The compounds represented by structure I have been shown to exhibit a variety of antifungal and antiparasitic activities to various degrees. For example, the compounds of structure I can inhibit the growth of various infectious fungi including Candida spp. (e.g., C. albicans, C. parapsilosis, C. krusei, C. glabrata, C. tropicalis, or C. lusitaniae), Torulopus spp. (e.g., T. glabrata), Aspergillus spp. (e.g., A. fumigatus), Histoplasma spp. (e.g., H. capsulatum), Cryptococcus spp. (e.g., C. neoformans), Blastomyces spp. (e.g., B. dermatitidis), Fusarium spp., Trichophyton spp., Pseudallescheria boydii, Coccidioides immitis, Sporothrix schenckii and the like.

Antifungal activity of a test compound is determined *in vitro* by obtaining the minimum inhibitory concentration (MIC) of the compound using a standard agar dilution test or a disc-diffusion test. The compound is then tested *in vivo* (in mice) to determine the effective dose for controlling a systemic fungal infection.

5

Accordingly, representative compounds of the present invention were tested for, and displayed, antifungal activity against at least one of the following fungii: C. albicans, C. parapsilosis, C. neoformans, Histoplasma spp, and A. fumigatus.

The compounds also inhibit growth of certain organisms primarily responsible for opportunistic infections in immunosuppressed individuals. For example, the compounds inhibit the growth of *Pneumocystis carinii* the causative organism of pneumocystis pneumonia (PCP) in AIDS and other immunocompromised recipients. "Topley and Wilson's Microbiology and Microbial Infections," Vol. 5, Ch. 22, Oxford University Press, Inc., New York, N.Y., 1998. Other protozoans that are inhibited by compounds of formula I include *Plasmodium spp.*, *Leishmania spp.*, *Trypanosoma spp.*, *Cryptosporidium spp.*, *Isospora spp.*, *Cyclospora spp.*,

Trichomonas spp., Microsporidiosis spp. and the like.

15

20

10

The dose of the compound represented by structure I administered varies depending on such factors as the nature and severity of the infection, the age and general health of the recipient and the tolerance of the recipient to the active ingredient. The particular dose regimen likewise can vary according to such factors and can be given in a single daily dose or in multiple doses during the day. The regimen can last from about 2 - 3 days to about 2 - 3 weeks or longer. A typical daily dose (administered in single or divided doses) contains a dosage level of from about 0.01 mg/kg to about 100 mg/kg of body weight of the active compound. Preferred daily doses are generally from about 0.1 mg/kg to about 60 mg/kg, more preferably from about 2.5 mg/kg to about 40 mg/kg.

25

Compound I can be administered parenterally, for example using intramuscular, sub-cutaneous, or intra-peritoneal injection, nasal, or oral means. In addition to these methods of administration, compound I can be applied topically for skin infections.

30

The invention also provides pharmaceutical formulations useful for administering the compounds of the invention. The active ingredient in such formulations comprises from 0.1% to 99.9% by weight of the formulation, more generally from about 10% to about 30% by weight.

For parenteral administration, the formulation comprises Compound I and a physiologically acceptable diluent such as deionized water, physiological saline, 5% dextrose and other commonly used diluents. The formulation can contain a solubilizing agent such as a polyethylene glycol or polypropylene glycol or other known solubilizing agent. Such formulations can be made up in sterile vials containing the active ingredient and one or more excipients in a dry powder or lyophilized powder form. Prior to use, a physiologically acceptable diluent is added and the solution withdrawn via syringe for administration to the recipient.

5

10

15

20

25

30

The pharmaceutical formulations are prepared by known procedures using known and readily available ingredients. In making the compositions of the present invention, the active ingredient will generally be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which can be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it can be a solid, semisolid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active ingredient, soft and hard gelatin capsules, suppositories, sterile injectable solutions, sterile packaged powders and the like.

For oral administration, the active ingredient is filled into gelatin capsules or formed into tablets. Tablets can also contain a binding agent, a dispersant or other excipients suitable for preparing a proper size tablet for the dosage and particular compound of the formula I. For pediatric or geriatric use the active ingredient can be formulated into a flavored liquid suspension, solution or emulsion. A preferred oral formulation is linoleic acid, cremophor RH-60 and water and preferably in the amount (by volume) of 8% linoleic acid, 5% cremophor RH-60, 87% sterile water and a compound of formula I in an amount of from about 2.5 to about 40 mg/mL.

For topical use the active ingredient can be formulated with a dry powder for application to the skin formulated in a liquid formulation comprising a solubilizing aqueous liquid or non-aqueous liquid, e.g., an alcohol or glycol.

#### **Formulations**

The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way. The term "active ingredient" refers to a compound represented by structure I or a pharmaceutically acceptable salt thereof.

Formulation Example 1

5

10

30

Hard gelatin capsules are prepared using the following ingredients:

	(mg/capsule)
Active ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

### Formulation Example 2

A tablet is prepared using the ingredients below:

15	•	Quantity (mg/capsule)
•	Active ingredient	250
	Cellulose, microcrystalline	400
	Silicon dioxide, fumed	10
20	Stearic acid	<u>5</u>
	Total	665 mg

The components are blended and compressed to form tablets of 665 mg each.

### Formulation Example 3

An aerosol solution is prepared containing the following components:

25		Weight
	Active ingredient	0.25
	Ethanol	25.75
	Propellant 22 (Chlorodifluoromethane) Total	74.00 100.00

The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

## Formulation Example 4

Tablets, each containing 60 mg of active ingredient, are made as follows:

	Active ingredient	60 mg
	Starch	45 mg
	Microcrystalline cellulose	35 mg
5	Polyvinylpyrrolidone (as 10% solution in water) Sodium carboxymethyl starch	4 mg 4.5 mg
	Magnesium stearate	0.5 mg
	Talc	<u>1 mg</u>
	Total	150 mg

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinyl-pyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

## Formulation Example 5

Capsules, each containing 80 mg of active ingredient, are made as follows:

	Active ingredient	80 mg
20	Starch	59 mg
	Microcrystalline cellulose	59 mg
	Magnesium stearate	<u>2 mg</u>
	Total	200 mg

10

15

25

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

## Formulation Example 6

Suppositories, each containing 225 mg of active ingredient, are made as follows:

30		Active ingredient	225 mg
		Saturated fatty acid glycerides	2,000 mg
	•	Total	2,225 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum

heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

### Formulation Example 7

Suspensions, each containing 50 mg of active ingredient per 5 mL dose, are made as follows:

Active ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mL
Benzoic acid solution	0.10 mL
Flavor	q.v.
Color	q.v.
Purified water to total	5 mL

5

10

15

20

The active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

## Formulation Example 8

An intravenous formulation can be prepared as follows:

Active ingredient 100 mg

Isotonic saline 1,000 mL

The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 mL per minute.

#### **CLAIMS**

WE CLAIM:

1. A compound represented by structure I:

5

10

wherein

R is an alkyl group, an alkenyl group, an alkynyl group, an aryl group, or heteroaryl group;

R<sup>1</sup> is independently -H, -OH or -O-Pg;

 $R^2$  is -H, -CH<sub>3</sub>, -NH<sub>2</sub>, or -NH-Pg;

R<sup>3</sup> is -H, -CH<sub>3</sub>, -CH<sub>2</sub>CONH<sub>2</sub>, -CH<sub>2</sub>CONH-Pg, -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, or -CH<sub>2</sub>CH<sub>2</sub>NH-Pg;

R<sup>4</sup> is -H, -OH, or -O-Pg;

R<sup>5</sup> is -OH, -OSO<sub>3</sub>H, or -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl,

15 C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, p-halophenyl, p-halophenoxy, p-nitrophenyl, p-nitrophenoxy, benzyl, benzyloxy, p-halobenzyl, p-halobenzyloxy, p-nitrobenzyl, or p-nitrobenzyloxy;

R<sup>6</sup> is -H, -OH, or -OSO<sub>3</sub>H;

R<sup>7</sup> is -H or -CH<sub>3</sub>;

t is an integer from 2-7;

R<sup>8</sup> is a sugar moiety of the formula

where R<sup>9</sup> is independently -H, -OH, -N<sub>3</sub>, -O-Pg, -NH<sub>2</sub>, -NH-Pg, or a second sugar moiety comprising one to three sugar units selected from the group consisting of

$$R^{9b}$$
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 

$$R^{9h} \xrightarrow{R^{9a}} R^{9a}$$

$$R^{9a} \xrightarrow{R^{9a}} R^{9a}$$

and mixtures

5 thereof, wherein

15

R<sup>9a</sup> is -H, -OH, -N<sub>3</sub>, -NH<sub>2</sub>, -O-Pg, or -NH-Pg,
R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or -NH-Pg,
R<sup>9c</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>N<sub>3</sub>, -CH<sub>2</sub>OSO<sub>3</sub>H, -CH<sub>2</sub>NH-Pg,
-CH<sub>2</sub>O-Pg, -CO<sub>2</sub>H, or -CO<sub>2</sub>-Pg,

10 where Ra is as defined above,

and so long as no more than one R9 is represented by said second sugar moiety;

Pg is a protecting group; and

pharmaceutically acceptable salts, esters, hydrates or solvates thereof.

2. The compound of Claim 1 wherein R is

where A, B, C and D are independently hydrogen, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>1</sub>-C<sub>12</sub> alkoxy, C<sub>1</sub>-C<sub>12</sub> alkylthio, halo, or
-O-(CH<sub>2</sub>)<sub>m</sub>-[O-(CH<sub>2</sub>)<sub>n</sub>]<sub>p</sub>-O-(C<sub>1</sub>-C<sub>12</sub> alkyl) or -O-(CH<sub>2</sub>)<sub>q</sub>-X-E;

m is 2, 3 or 4; n is 2, 3 or 4; p is 0 or 1; q is 2, 3 or 4;
X is pyrrolidino, piperidino or piperazino; and E is hydrogen, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, benzyl or C<sub>3</sub>-C<sub>12</sub> cycloalkylmethyl.

3. The compound of Claim 2 wherein

R<sup>1</sup> is hydroxy at each occurrence;

R<sup>4</sup> is hydroxy;

10

15

20

R<sup>5</sup> is -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is methyl or methoxy;

 $R^2$ ,  $R^3$ , and  $R^7$  are each methyl; and

R is a moiety of the formula

a pharmaceutically acceptable salt or solvate thereof.

4. The compound of Claim 3 wherein

R<sup>5</sup> is hydroxy;

R is a moiety of the formula

where C is hydrogen or C3-C7 alkoxy;

R<sup>8</sup> is a moiety of the formula

where R<sup>9</sup> is independently hydrogen, hydroxy, amino. or a moiety of the formula

5

 $R^{9b}$  is -OPO<sub>2</sub> $R^a$ , -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or

-NH-Pg and n is 1, 2, or 3; or

a pharmaceutically acceptable salt or solvate thereof.

10

5. The compound of Claim 4 wherein

C is n-pentoxy;

R<sup>9</sup> is independently hydroxy or amino; or a pharmaceutically acceptable salt or solvate thereof.

15

25

6. The compound of claim 5 wherein

R9 is hydroxy at each occurrence; and

t is 2; or a pharmaceutically acceptable salt of solvate thereof.

- 7. A pharmaceutical formulation comprising one or more pharmaceutical carriers, diluents, or excipients and a compound of Claim 1.
  - 8. A method of inhibiting fungal activity comprising administering to a recipient in need of such inhibition an effective amount of a compound represented by structure I:

wherein

5

10

15

R is an alkyl group, an alkenyl group, an alkynyl group, an aryl group, or heteroaryl group;

R<sup>1</sup> is independently -H, -OH or -O-Pg;

R<sup>2</sup> is -H, -CH<sub>3</sub>, -NH<sub>2</sub>, or -NH-Pg;

 $R^3$  is -H, -CH<sub>3</sub>, -CH<sub>2</sub>CONH<sub>2</sub>, -CH<sub>2</sub>CONH-Pg, -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, or -CH<sub>2</sub>CH<sub>2</sub>NH-Pg;

R<sup>4</sup> is -H, -OH, or -O-Pg;

R<sup>5</sup> is -OH, -OSO<sub>3</sub>H, or -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, p-halophenyl, p-halophenoxy, p-nitrophenoxy, benzyl, benzyloxy, p-halobenzyl, p-halobenzyloxy, p-nitrobenzyl, or p-nitrobenzyloxy;

R<sup>6</sup> is -H, -OH, or -OSO<sub>3</sub>H;

R<sup>7</sup> is -H or -CH<sub>3</sub>;

t is an integer from 2-7;

R<sup>8</sup> is a sugar moiety of the formula

where R9 is independently -H, -OH, -N3, -O-Pg, -NH2,

-NH-Pg, or a second sugar moiety comprising one to three sugar units selected from the group consisting of

$$R^{9b} \longrightarrow R^{9a} \longrightarrow R$$

' and mixtures

thereof, wherein

5 R<sup>9a</sup> is -H, -OH, -N<sub>3</sub>, -NH<sub>2</sub>, -O-Pg, or -NH-Pg, R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or -NH-Pg, R<sup>9c</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>N<sub>3</sub>, -CH<sub>2</sub>OSO<sub>3</sub>H, -CH<sub>2</sub>NH-Pg,

-CH<sub>2</sub>O-Pg, -CO<sub>2</sub>H, or -CO<sub>2</sub>-Pg,

where Ra is as defined above,

and so long as no more than one R<sup>9</sup> is represented by said second sugar moiety;

Pg is a protecting group; and

pharmaceutically acceptable salts, esters, hydrates or solvates thereof.

9. The Method of Claim 8 wherein R is

$$\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \\ \end{array} \end{array} \end{array} \begin{array}{c} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \end{array} \begin{array}{c} \\ \end{array} \begin{array}{c} \\$$

where A, B, C and D are independently hydrogen, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkynyl,

C<sub>1</sub>-C<sub>12</sub> alkoxy, C<sub>1</sub>-C<sub>12</sub> alkylthio, halo, or

 $-O-(CH_2)_m-[O-(CH_2)_n]_p-O-(C_1-C_{12} \text{ alkyl}) \text{ or } -O-(CH_2)_q-X-E;$ 

m is 2, 3 or 4; n is 2, 3 or 4; p is 0 or 1; q is 2, 3 or 4;

5 X is pyrrolidino, piperidino or piperazino; and E is hydrogen, C1-C12 alkyl,

 $C_3$ - $C_{12}$  cycloalkyl, benzyl or  $C_3$ - $C_{12}$  cycloalkylmethyl.

10. The method of claim 8 wherein the recipient is a human.

10 11. The method of Claim 9 wherein

R1 is hydroxy at each occurrence;

R4 is hydroxy;

R<sup>5</sup> is -OPO<sub>2</sub>HR<sup>2</sup>, where R<sup>2</sup> is methyl or methoxy;

R<sup>2</sup>, R<sup>3</sup>, and R<sup>7</sup> are each methyl; and

R is a moiety of the formula

a pharmaceutically acceptable salt or solvate thereof.

12. The method of Claim 9 wherein

R<sup>5</sup> is hydroxy;

R is a moiety of the formula

where C is hydrogen or C3-C7 alkoxy;

R<sup>8</sup> is a moiety of the formula

25

where  $R^{9}$  is independently hydrogen, hydroxy, amino. or a moiety of the formula

R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or

5 -NH-Pg and n is 1, 2, or 3; or

a pharmaceutically acceptable salt or solvate thereof.

13. The method of Claim 12 wherein

C is n-pentoxy;

10 R<sup>9</sup> is independently hydroxy or amino; or a pharmaceutically acceptable salt or solvate thereof.

14. The method of claim 13 wherein

R9 is hydroxy at each occurrence; and

t is 2; or a pharmaceutically acceptable salt of solvate thereof.

15. The method according to Claim 8 wherein the fungal activity arises from one or more of the fungi selected from the group consisting of *Candida albicans*, *Aspergillus fumigatis*, and *Candida parapsilosis*.

20

15

16. A method of inhibiting parasitic activity comprising administering to a recipient in need of such inhibition an effective amount of a compound represented by structure I:

wherein

5

10

15

R is an alkyl group, an alkenyl group, an alkynyl group, an aryl group, or heteroaryl group;

R<sup>1</sup> is independently -H, -OH or -O-Pg;

R<sup>2</sup> is -H, -CH<sub>3</sub>, -NH<sub>2</sub>, or -NH-Pg;

 $R^3$  is -H, -CH $_2$ , -CH $_2$ CONH $_2$ , -CH $_2$ CONH-Pg, -CH $_2$ CH $_2$ NH-Pg;

R<sup>4</sup> is -H, -OH, or -O-Pg;

R<sup>5</sup> is -OH, -OSO<sub>3</sub>H, or -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is hydroxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, p-halophenyl, p-halophenoxy, p-nitrophenyl, p-nitrophenoxy, benzyl, benzyloxy, p-halobenzyl, p-halobenzyloxy, p-nitrobenzyl, or p-nitrobenzyloxy;

R<sup>6</sup> is -H, -OH, or -OSO<sub>3</sub>H;

R<sup>7</sup> is -H or -CH<sub>3</sub>;

t is an integer from 2-7;

R<sup>8</sup> is a sugar moiety of the formula

$$R^9$$
  $R^9$   $R^9$   $R^9$   $R^9$ 

where R9 is independently -H, -OH, -N3, -O-Pg, -NH2,

-NH-Pg, or a second sugar moiety comprising one to three sugar units selected from the group consisting of

$$R^{9b}$$
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 
 $R^{9a}$ 

and mixtures

thereof, wherein

R<sup>9a</sup> is -H, -OH, -N<sub>3</sub>, -NH<sub>2</sub>, -O-Pg, or -NH-Pg, 5 R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or -NH-Pg, R% is -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>N<sub>3</sub>, -CH<sub>2</sub>OSO<sub>3</sub>H, -CH<sub>2</sub>NH-Pg,

-CH<sub>2</sub>O-Pg, -CO<sub>2</sub>H, or -CO<sub>2</sub>-Pg,

where Ra is as defined above,

and so long as no more than one R9 is represented by said second sugar moiety; 10 Pg is a protecting group; and

pharmaceutically acceptable salts, esters, hydrates or solvates thereof.

#### 17. The Method of Claim 16 wherein R is

where A, B, C and D are independently hydrogen, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkynyl,

C<sub>1</sub>-C<sub>12</sub> alkoxy, C<sub>1</sub>-C<sub>12</sub> alkylthio, halo, or

-O-(CH<sub>2</sub>)<sub>m</sub>-[O-(CH<sub>2</sub>)<sub>n</sub>]<sub>p</sub>-O-(C<sub>1</sub>-C<sub>12</sub> alkyl) or -O-(CH<sub>2</sub>)<sub>q</sub>-X-E;

m is 2, 3 or 4; n is 2, 3 or 4; p is 0 or 1; q is 2, 3 or 4:

X is pyrrolidino, piperidino or piperazino; and E is hydrogen, C1-C12 alkyl,

C<sub>3</sub>-C<sub>12</sub> cycloalkyl, benzyl or C<sub>3</sub>-C<sub>12</sub> cycloalkylmethyl.

18. The method of claim 16 wherein the recipient is a human.

10 19. The method of Claim 17 wherein


R<sup>1</sup> is hydroxy at each occurrence;

R<sup>4</sup> is hydroxy;

R<sup>5</sup> is -OPO<sub>2</sub>HR<sup>a</sup>, where R<sup>a</sup> is methyl or methoxy;

R<sup>2</sup>, R<sup>3</sup>, and R<sup>7</sup> are each methyl; and

15 R is a moiety of the formula



a pharmaceutically acceptable salt or solvate thereof.

20. The method of Claim 17 wherein

20  $R^5$  is hydroxy;

R is a moiety of the formula

where C is hydrogen or C3-C7 alkoxy;

R<sup>8</sup> is a moiety of the formula

where  $R^{9}$  is independently hydrogen, hydroxy, amino. or a moiety of the formula

R<sup>9b</sup> is -OPO<sub>2</sub>R<sup>a</sup>, -OSO<sub>3</sub>H, -H, -NH<sub>2</sub>, -OH, -O-Pg, or

- 5 -NH-Pg and n is 1, 2, or 3; or
  - a pharmaceutically acceptable salt or solvate thereof.
  - 21. The method of Claim 20 wherein

C is n-pentoxy;

- 10 R<sup>9</sup> is independently hydroxy or amino; or a pharmaceutically acceptable salt or solvate thereof.
  - 22. The method of claim 21 wherein
  - R<sup>9</sup> is hydroxy at each occurrence; and
- t is 2; or a pharmaceutically acceptable salt of solvate thereof.
  - 23. The method of Claim 16 wherein the parasitic activity arises from *Pneumocystis carinii*.

Into. onal Application No PCT/US 99/29927

	PC1/US 99/2992/
A CLASSIFICATION OF SUBJECT MATTER IPC 7 C07K7/56 A61K38/12 A61P31/	10 A61P33/00
According to International Patent Classification (IPC) or to both national classific	cation and IPC
B. FIELDS SEARCHED	
Minimum documentation searched (classification system tollowed by classificat IPC 7 CO7K A61K	ion symbole)
Documentation searched other than minimum documentation to the extent that  Electronic data base consulted during the international search (name of data by	
	эээ анд, wixer рислем, воагси юнно цоочу
C. DOCUMENTS CONSIDERED TO BE RELEVANT	T Bright Hall
Category * Citation of document, with indication, where appropriate, of the re	levant passages Relevant to claim No.
Y EP 0 757 058 A (LILLY CO ELI) 5 February 1997 (1997-02-05) the whole document	1-23
P,Y WO 99 43337 A (LILLY CO ELI ;JAM ANDREW (US); RODRIGUEZ MICHAEL JO 2 September 1999 (1999-09-02) the whole document	
A WO 97 17365 A (MERCK & CO INC ;BA JAMES M (US); BOUFFARD FRANCES A 15 May 1997 (1997-05-15) the whole document	ALKOVEC (US); DR)
Further documents are listed in the continuation of box C.	Patent family members are listed in annex.
*Special categories of cited documents:  *A* document defining the general state of the art which is not considered to be of particular relevance  *E* earlier document but published on or after the international filling data  *I.* document which may throw doubts on priority ctaim(s) or which is cited to establish the publication date of enother citation or other special reason (as specified)  *O* document referring to an oral disclosure, use, exhibition or other means  *P* document published prior to the international filing date but later than the priority date claimed  Date of the actual completion of the international search  9 May 2000	To later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combined with one or more other such documents, such combination being obvious to a person sidiled in the art.  "&" document member of the same patent family  Date of mailing of the International search report
Name and mailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (431-70) 340-2040, Tx. 31 651 epo nl, Fax: (431-70) 340-3016	Authorized officer  Groenendijk, M

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

Inamational application No.

PCT/US 99/29927

BxI	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This inte	emational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claims 8-23 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful international Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This inte	mational Searching Authority found multiple inventions in this international application, as follows:
. —	
'· 📋	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest  The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

information on patent family members

Into. onal Application No PCT/US 99/29927

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP 0757058	A	05-02-1997	US	5629289 A	13-05-1997
			AU	6593896 A	26-02-1 <b>997</b>
			JP	11510165 T	07 <b>-</b> 09-1 <b>99</b> 9
			WO	9705163 A	13-02-1 <b>997</b>
WO 9943337	Α	02-09-1999	AU	2686099 A	15-09-1 <b>99</b> 9
			EP	0943623 A	22-09-1 <b>99</b> 9
WO 9717365	Α	15-05-1997	AU	707664 B	15-07-1 <b>99</b> 9
			AU	7554096 A	29 <b>-</b> 05-1 <b>99</b> 7
			EP	0862579 A	09-09-1 <b>99</b> 8
			HR	960521 A	30-04-1 <b>99</b> 8
			JP 2	000501076 T	02-02-2000